

acts as a liver protector
and purifies the body

Animal Nutrition
Animal Health

SANILIVER PLUS



PROTECTION AGAINST LIVER DAMAGE

IMPROVED DIGESTIVE FUNCTION

IMPROVING PRODUCTIVITY



SANILIVER PLUS

SANILIVER PLUS provides support to the liver, so that it accelerates its recovery in periods of intense activity or overload, improving production.

SANILIVER PLUS is a food that contains several substances with hepatoprotective properties, that is, they are responsible for protecting hepatocytes (liver cells) against toxic agents. Hepatotropic agents are inserted into the group of hepatoprotectors, that is, they have a special tropism for the liver, and also lipotropic agents, which lead to the elimination of lipids or decreased deposition in the liver.

SANILIVER PLUS, among others, contains vitamin E, vitamin C, vitamin B12, methionine and choline.

Vitamin E and Vitamin C: These two agents are believed to have an antioxidant effect and therefore have an antinecrotic effect;

Vitamin B12: is a lipotropic agent that favors the synthesis of liver proteins. This vitamin is also involved in the formation of choline and in the biotransformation of labile methyl radicals;

Methionine: it is a donor of the methyl radical, which favors the methylation of different chemical substances, which facilitates its elimination from the body. Methionine donates methyl radical even for choline synthesis. Methionine also appears to have an antinecrotic effect on the liver;

Choline: This agent is considered lipotropic. It is believed to be capable of promoting the conversion of liver fat to choline-containing phospholipids, which can be rapidly transferred from the liver into the bloodstream, thereby preventing liver stenosis.

I

INDICATIONS

SANILIVER PLUS is indicated for fatty liver, infections and liver problems, food withholdings, enterotoxemias and intoxications, mycotoxicosis and medications.

HOW TO USE

Orally, in drinking water, milk substitutes or in liquid foods, according with the indications expressed in the instructions for use.

DOSAGE

1 to 2 ml. per liter

for 5 to 7 days, for: chicken (1st and 4th weeks); field chickens (1st, 4th and 9th weeks); turkeys (1st, 4th, 8th and 12th weeks); Ducks (1st and 5th weeks); quail (1st and 3rd weeks); recreational birds (1st and 5th weeks); breeders and lay (start of lay, sunset peak and 45th week); pigs and rabbits

10 to 20 ml. per liter

for 5 to 7 days for Calves and Pigs (boars and young)

2 ml. for 50 kg of weight

body for 5 to 7 days for Sheep.



C/ Gutemberg, 61 P.I. Villares - 37184 Villares de la Reina - Salamanca - SPAIN

Tel. 0034 923 234 866 0034 * 923 600 090 - salpesa@salpesa.com - www.salpesa.com